

Product Name: CANIGEN DHA2P TRIVALENT LIVING VACCINE
APVMA Approval No: 40758/115156



Label Name:	CANIGEN DHA2P TRIVALENT LIVING VACCINE
Signal Headings:	PRESCRIPTION ANIMAL REMEDY KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY
Constituent Statements:	Each single dose vial contains (minimum titres in TCID50): <ul style="list-style-type: none">- live, attenuated canine distemper virus 10^3- live, attenuated canine adenovirus type 2 (infectious hepatitis) 10^4- live, attenuated canine parvovirus 10^5
Claims:	For vaccination of dogs against canine distemper virus (CDV), canine adenovirus (CAV) (infectious hepatitis) and canine parvovirus (CPV) and the reduction of shedding of CPV. This vaccine provides cross-protection against CPV-2, CPV-2b and CPV-2c.
Net Contents:	10 x 1mL single dose vials. 20 x 1mL single dose vials.
Directions for Use:	
Restrains:	
Contraindications:	
Precautions:	Do not vaccinate dogs with a compromised immune status or receiving immunosuppressive drugs. Do not vaccinate pregnant bitches.
Side Effects:	Adverse post-vaccination reactions such as anaphylaxis/hypersensitivity occur very rarely and may require appropriate treatment.

Dosage and Administration:	<p>Only healthy dogs with normal temperatures should be vaccinated after a clinical examination.</p> <p>Do not use chemicals to disinfect the skin, needles or syringes before vaccination as this could inactivate the viruses.</p> <p>Use immediately after reconstitution with 1 mL of a suitable diluent (i.e. water for injections BP or Canigen KC Bivalent Inactivated Vaccine for Dogs) transferred aseptically.</p> <p>Vaccination Programs: Primary vaccination in puppies from 8 weeks of age: Inject subcutaneously 1 dose of vaccine at 8 weeks and 1 dose at 12 weeks. Under normal circumstances, 2 doses are sufficient to ensure protective immunisation in puppies from 8 weeks of age.</p> <p>Primary vaccination in dogs over 12 weeks of age: Inject subcutaneously 1 dose of vaccine. Under normal circumstances 1 dose of vaccine will be sufficient to ensure protective immunisation in puppies at or over 12 weeks of age.</p> <p>In situations where high maternal antibody and potential challenge is possible an additional vaccination should be considered at 16 weeks of age.</p> <p>Primary vaccination in adult dogs: Inject subcutaneously 1 dose of vaccine.</p> <p>One dose of vaccine should be given 12 months after the initial primary vaccination.</p> <p>Revaccination: A single dose vaccine should be given at intervals of up to 3 years following the primary vaccination program.</p> <p>Many factors influence the effectiveness of vaccination and the need for revaccination. Taking all these factors into account, each individual animal should be vaccinated as frequently as considered necessary by their veterinarian to provide protection.</p>
----------------------------	--

General Directions:	
---------------------	--

Withholding Periods:	
----------------------	--

Trade Advice:	
---------------	--

Safety Directions:	
--------------------	--

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre: Phone Australia 131126.
-------------------------	--

First Aid Warnings:	
---------------------	--

Additional User Safety:	Take Care to Avoid Self-Injection. In the event of self-administration, seek medical attention if you are concerned. Show the package leaflet or label to the Medical Practitioner.
-------------------------	---

Environmental Statements:	
---------------------------	--

Disposal:	Dispose of empty containers and outer packaging by wrapping with paper and putting in garbage. Discarded needles/sharps should immediately be placed in a designated and appropriately labelled "sharps" container. Discarded vials should immediately be placed in a designated and appropriately labelled "biologicals" container.
-----------	---

Storage:	Store between 2°C and 8°C (refrigerate, do not freeze). Protect from light.
----------	---