



Product Name: MASTALONE INTRAMAMMARY SUSPENSION FOR LACTATING COWS  
APVMA Approval No: 49851/106743

Label Name:	MASTALONE INTRAMAMMARY SUSPENSION FOR LACTATING COWS
-------------	--

Signal Headings:	PRESCRIPTION ANIMAL REMEDY KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY
------------------	--

Constituent Statements:	185 mg/10ml OXYTETRACYCLINE AS OXYTETRACYCLINE HYDROCHLORIDE 100 mg/10ml OLEANDOMYCIN AS PHOSPHATE 100 mg/10ml NEOMYCIN AS SULFATE
-------------------------	--

Claims:	A broad spectrum antibiotic combination for the treatment of mastitis in lactating cattle
---------	---

Net Contents:	20 x 10mL
---------------	-----------

Directions for Use:	
---------------------	--

Dosage and Administration:	<p>Infuse 1 syringe every 24 hours for 3 treatments per infected quarter. In cold weather warm syringe to body temperature.</p> <p>The entire contents (10 mL) of one syringe should be infused into the affected quarter immediately after milking. Ensure that the affected quarters are thoroughly stripped out before treatment. Treatment should be repeated at 24 hour intervals for three days. Care should be taken to properly clean and disinfect the tip of the teat to prevent introduction of contaminants. Care should also be taken to avoid contamination of the tip of the syringe after removal of the cap and after prior to insertion into the teat canal. After contents are infused, gently massage the preparation upwards, working the syringe contents into the mammary gland.</p> <p>In addition to therapeutic treatment, cleanliness of dairy, milking equipment, thoroughness of milking techniques, operator care and correctly adjusted milking machines are of paramount importance.</p> <p>The routine use of suitable commercial teat dips for all lactating cows is highly recommended.</p>
----------------------------	--

General Directions:	The combination of three antibiotics in MASTALONE is designed to provide maximum coverage against most of the known causative organisms associated with bacterial mastitis.
---------------------	---

	<p>Terramycin* (oxytetracycline hydrochloride) is active against wide range of Gram positive and Gram negative organisms.</p> <p>Oleandomycin provides specific reinforcement against Gram positive organisms including staphylococci and streptococci which may be resistant to penicillin and streptomycin.</p> <p>Neomycin provides specific reinforcement against Gram negative organisms including E. coli, Proteus and Pseudomonas.</p>
--	---

Withholding Periods:	<p><b>WITHHOLDING PERIODS</b></p> <p>MILK: milk taken from animals within 7 days following treatment <b>MUST NOT BE USED</b> for human consumption or supplied for processing.</p> <p>MEAT: <b>DO NOT USE</b> less than 30 days before slaughter for human consumption.</p>
----------------------	---

Trade Advice:	<p><b>EXPORT SLAUGHTER INTERVAL (ESI):</b> This product does not have an ESI established. For advice on the ESI, contact the manufacturer on 1800 884 813 before using this product.</p>
---------------	--

First Aid Instructions:	<p><b>FIRST AID</b></p> <p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.</p>
-------------------------	---

First Aid Warnings:	
---------------------	--

Disposal:	<p><b>DISPOSAL</b></p> <p>Dispose of empty syringes by wrapping with paper and putting in garbage.</p>
-----------	--

Storage:	<p><b>STORAGE</b></p> <p>Store below 25°C (Air-Conditioning)</p>
----------	--