

Main panel – draft immediate label



**PRESCRIPTION ANIMAL REMEDY  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY**



**ILIUM ACEPRIL-10  
INJECTION  
TRANQUILLISER, PRE-ANAESTHETIC MEDICATION**

ACTIVE CONSTITUENT:  
10 mg/mL ACEPROMAZINE (as MALEATE)

ILIUM 20mL

Ancillary panels

Tranquilliser, Pre-anaesthetic medication for horses, cattle, sheep and pigs.

**DIRECTIONS FOR USE:**

**Dosage and Administration**

Use contents within 24 hours of first broaching of the vial. Discard the unused portion.

**HORSES, CATTLE, SHEEP & PIGS:**

0.25-0.5 mL per 50 kg bodyweight i.m. or slow i.v.

**WITHHOLDING PERIODS:**

**MEAT (CATTLE, SHEEP AND PIGS): DO NOT USE less than 2 days before slaughter for human consumption.**

**MEAT (HORSES): DO NOT USE less than 28 days before slaughter for human consumption.**

**MILK (COWS AND EWES): Milk collected from cows or ewes within 48 hours (4 milkings) following treatment MUST NOT BE USED for human consumption or processing, or fed to bobby calves.**

**TRADE ADVICE:**

EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact the manufacturer on (02) 9604 6266 before using this product.

**First Aid:**

If poisoning occurs, contact a doctor or Poisons Information Centre.

Phone Australia 131126

TROY LABORATORIES PTY. LIMITED

31-35 Glendenning Road, Glendenning NSW 2761

website: [www.troylab.com.au](http://www.troylab.com.au) email: [info@troylab.com.au](mailto:info@troylab.com.au)

Dispose of empty container by wrapping with paper and putting in the garbage.

Store below 25°C (Air Conditioning). Protect from light.

APVMA Approval No. 50521/50346

Batch:.

Expiry:

*Main panel – draft immediate label*



**PRESCRIPTION ANIMAL REMEDY  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY**

**ILIUM ACEPRIL-10  
INJECTION  
TRANQUILLISER, PRE-ANAESTHETIC MEDICATION**

ACTIVE CONSTITUENT:  
10 mg/mL ACEPROMAZINE (as MALEATE)

ILIUM 20mL

*Ancillary  
panels*

Tranquilliser, Pre-anaesthetic medication for horses, cattle, sheep and pigs.

**DIRECTIONS FOR USE:**

**Dosage and Administration**

Use contents within 24 hours of first broaching of the vial. Discard the unused portion.

**HORSES, CATTLE, SHEEP & PIGS:**

0.25-0.5 mL per 50 kg bodyweight i.m. or slow i.v.

**WITHHOLDING PERIODS:**

**MEAT (CATTLE, SHEEP AND PIGS): 2 days**

**MEAT (HORSES): 28 days**

**MILK (COWS AND EWES): 2 days**

**TRADE ADVICE:** EXPORT SLAUGHTER INTERVAL (ESI): ESI Not established.

**First Aid:** If poisoning occurs, contact a doctor or Poisons Information Centre.  
*Phone Australia 131126*

TROY LABORATORIES PTY. LTD  
31-35 Glendenning Road, Glendenning NSW 2761

Dispose of empty container by wrapping with paper and putting in the garbage.

Store below 25°C (Air Conditioning). Protect from light.

APVMA Approval No. 50521/50346

Batch:

Expiry: