

Main panel – draft label

**PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**



**BENACILLIN
LONG ACTING PENICILLIN INJECTION**



ACTIVE CONSTITUENT:
150 mg/mL PROCAINE PENICILLIN
150 mg/mL BENZATHINE PENICILLIN
20 mg/mL PROCAINE HYDROCHLORIDE

For parenteral treatment of infections caused by bacteria sensitive to penicillin
in dogs, cats, pigs, calves, foals, horses and cattle

ILIUM 100mL/250mL

Ancillary panel – draft label

**DIRECTIONS FOR USE:
SHAKE WELL BEFORE USE
Dosage and Administration**

Use contents within 24 hours of first broaching of the vial. Discard the unused portion.

By i.m. administration only

DOGS & CATS:	0.5 - 4 mL	CALVES & FOALS:	3 - 7 mL
PIGS:	3 - 7 mL	HORSES & CATTLE:	10 - 20 mL

Therapeutic levels maintained for up to 5 days.

WITHHOLDING PERIODS:

MILK: Milk collected from cows within 13 days following treatment
MUST NOT BE USED for human consumption or processing. This milk
should not be fed to bobby calves.

MEAT :CATTLE, PIGS AND HORSES: DO NOT USE less than 30 days
before slaughter for human consumption.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI
established. For advice on the ESI, contact the manufacturer on (02) 9604
6266 before using this product.

First Aid:

If poisoning occurs, contact a doctor or Poisons Information Centre.
Phone Australia 131126

TROY LABORATORIES PTY. LTD
31-35 Glendenning Road, Glendenning NSW 2761

website: www.troylab.com.au email: info@troylab.com.au

Dispose of empty container by wrapping with paper and putting in the garbage.

Store below 25°C (Air Conditioning). Protect from light.

APVMA Approval No. 50522/50297

Batch:.

Expiry:

Main panel – draft label

**PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**



**BENACILLIN
LONG ACTING PENICILLIN INJECTION**

ACTIVE CONSTITUENT:
150 mg/mL PROCAINE PENICILLIN
150 mg/mL BENZATHINE PENICILLIN
20 mg/mL PROCAINE HYDROCHLORIDE

For parenteral treatment of infections caused by bacteria sensitive to penicillin
in dogs, cats, pigs, calves, foals, horses and cattle

ILIUM 100mL/250mL

Ancillary panel – draft label

**DIRECTIONS FOR USE:
SHAKE WELL BEFORE USE
Dosage and Administration**

Use contents within 24 hours of first broaching of the vial. Discard the unused portion.

By i.m. administration only

DOGS & CATS:	0.5 - 4 mL	CALVES & FOALS:	3 - 7 mL
PIGS:	3 - 7 mL	HORSES & CATTLE:	10 - 20 mL

Therapeutic levels maintained for up to 5 days.

WITHHOLDING PERIODS:

MILK: Milk collected from cows within 13 days following treatment
MUST NOT BE USED for human consumption or processing. This milk
should not be fed to bobby calves.

MEAT :CATTLE, PIGS AND HORSES: DO NOT USE less than 30 days
before slaughter for human consumption.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI
established. For advice on the ESI, contact the manufacturer on (02) 9604
6266 before using this product.

First Aid:

If poisoning occurs, contact a doctor or Poisons Information Centre.
Phone Australia 131126

TROY LABORATORIES PTY. LTD

31-35 Glendenning Road, Glendenning NSW 2761
website: www.troylab.com.au email: info@troylab.com.au

Dispose of empty container by wrapping with paper and putting in the garbage.

Store below 25°C (Air Conditioning). Protect from light.

APVMA Approval No. 50522/50297

Batch:.

Expiry: