

**PRIMARY PACK - OUTER CARTON LABEL****MAIN PANEL**

**PRESCRIPTION ANIMAL REMEDY  
KEEP OUT OF THE REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY**

**OPTIMMUNE ®OPHTHALMIC OINTMENT**

**ACTIVE CONSTITUENT:** 2 mg/g CYCLOSPORIN

For the treatment of idiopathic chronic keratoconjunctivitis sicca (KCS) and chronic superficial keratitis (CSK) in dogs.

**6 x 3.5 g Tubes**

**PRIMARY PACK – OUTER CARTON LABEL****ANCILLIARY PANELS****DIRECTIONS FOR USE****Dosage and administration**

Remove debris from affected eye(s) with suitable non-irritating solution.

For KCS apply a 6 mm strip of ointment to the affected eye(s) every 12 hours.

For CSK apply a 6-10 mm strip of ointment to the affected eye(s) every 12 hours.

Place the ointment directly on the cornea or into the conjunctival sac. Continue treatment while condition persists. Dogs afflicted with KCS or CSK may require lifelong consistent therapy.

**General directions**

For best results in treating KCS the ointment should be administered early in the course of the disease before irreversible damage to the lacrymal tissue, or dense corneal scarring or pigmentation occurs. It is recommended that dogs exhibiting chronic recurring conjunctivitis be tested for adequate tear production.

The clinical effect of OPTIMMUNE Ophthalmic Ointment has not been determined in dogs with KCS due to the following conditions: congenital alacrima, sulfonamide usage, canine distemper virus, metabolic disease, surgical removal of the third eyelid gland and facial nerve paralysis with loss of the palpebral reflex. Several days to a few weeks of application may be required before the clinical effects of Optimimmune are of sufficient magnitude to allow the safe withdrawal of previously initiated therapy.

Abrupt withdrawal of a therapeutic agent immediately upon initiation of treatment with Optimimmune can result in rapid clinical relapse.

**Precautions/contraindications**

Do not use if pre-existing viral or fungal ocular infections are present. Delay use until such infections have been successfully treated. The safety of Optimimmune in puppies, pregnant bitches, or dogs used for breeding has not been determined.

On rare occasions, instillation of OPTIMMUNE may be associated with stinging. If this persists beyond 7 days treatment should be withdrawn.

Additional information is in the Material Safety Data Sheet.

**Dispose** of empty container by wrapping with paper and putting in garbage.

**Store** below 25°C (Air Conditioning).

Schering-Plough Animal Health  
(A Division of Schering-Plough Pty Ltd)  
11 Gibbon Road, Baulkham Hills NSW 2153

NRA 51551/0600

**BATCH**

**EXPIRY**

**IMMEDIATE CONTAINER TUBE LABEL****MAIN PANEL**

**PRESCRIPTION ANIMAL REMEDY**  
**KEEP OUT OF REACH OF CHILDREN**  
**FOR ANIMAL TREATMENT ONLY**

**OPTIMMUNE ®OPHTHALMIC OINTMENT**

**ACTIVE CONSTITUENT:** 2 mg/g CYCLOSPORIN

For the treatment of idiopathic chronic keratoconjunctivitis sicca (KCS) and chronic superficial keratitis (CSK) in dogs.

**3.5 g**

**IMMEDIATE CONTAINER TUBE LABEL****ANCILLIARY PANELS****DIRECTIONS FOR USE**

**READ THE ENCLOSED LEAFLET BEFORE USING THIS PRODUCT.**

For KCS and CSK apply a 6 mm strip of ointment to the affected eye(s) every 12 hours.

**Dispose** of empty container by wrapping with paper and putting in garbage.

**Store** below 25°C (Air conditioning).

Schering-Plough Animal Health  
(A Division of Schering-Plough Pty Ltd)  
11 Gibbon Road, Baulkham Hills NSW 2153

NRA 51551/0600

**(B) See crimp of tube**

**EXP. See crimp of tube**

**TAMPER PROOF PACKET, ENCLOSURES TUBE****MAIN PANEL**

**PRESCRIPTION ANIMAL REMEDY  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY**

**OPTIMMUNE ®OPHTHALMIC OINTMENT****Active Constituent:** 2 mg/g CYCLOSPORIN

For the treatment of idiopathic chronic keratoconjunctivitis sicca (KCS) and chronic superficial keratitis (CSK) in dogs.

Contents: 3.5 g

**TAMPER PROOF PACKET, ENCLOSURES TUBE****ANCILLIARY PANELS**

Tamper Evident – Sealed in a pouch for the dog's protection

**Dosage and administration**

Remove debris from affected eye(s) with suitable non-irritating solution.

For KCS and CSK apply a 6 mm strip of ointment to the affected eye(s) every 12 hours.

**DISPOSAL:** Dispose of empty container by wrapping with paper and putting in garbage.**STORAGE:** Store below 25°C (Air conditioning).

Schering-Plough Animal Health  
(A Division of Schering-Plough Pty Limited),  
11 Gibbon Road, Baulkham Hills NSW 2513

NRA 51551/0600

Expiry Date

Batch Number

**BOX FOR SINGLE TUBE****MAIN PANEL**

**PRESCRIPTION ANIMAL REMEDY  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY**

**OPTIMMUNE ®OPHTHALMIC OINTMENT**

**Active Constituent:** 2 mg/g CYCLOSPORIN

For the treatment of idiopathic chronic keratoconjunctivitis sicca (KCS) and chronic superficial keratitis (CSK) in dogs.

Contents: 3.5 g

**BOX FOR SINGLE TUBE****BACK PANEL****Dosage and administration**

Remove debris from affected eye(s) with suitable non-irritating solution.

For KCS apply a 6 mm strip of ointment to the affected eye(s) every 12 hours.

For CSK apply a 6-10 mm strip of ointment to the affected eye(s) every 12 hours.

Place the ointment directly on the cornea or into the conjunctival sac. Continue treatment while condition persists.

Dogs afflicted with KCS or CSK may require lifelong consistent therapy.

**DISPOSAL:** Dispose of empty container by wrapping with paper and putting in garbage.

**STORAGE:** Store below 25°C (Air conditioning).

Schering-Plough Animal Health  
(A Division of Schering-Plough Pty Limited),  
11 Gibbon Road, Baulkham Hills NSW 2513

NRA 51551/0600

**BOX FOR SINGLE TUBE****END PANEL 1****OPTIMMUNE ®OPHTHALMIC OINTMENT****BOX FOR SINGLE TUBE****END PANEL 2**

**Batch and Exp**

**BOX FOR SINGLE TUBE****SIDE PANEL 1****OPTIMMUNE ®OPHTHALMIC OINTMENT****Active Constituent:** 2 mg/g CYCLOSPORIN

® Schering Pough Pty Ltd 2000

© Schering Pough Pty Ltd 2000

**BOX FOR SINGLE TUBE****SIDE PANEL 2****OPTIMMUNE ®OPHTHALMIC OINTMENT****Directions** - For KCS and CSK apply a 6-10 mm strip of ointment to the affected eye(s) every 12 hours.

The 10mm measurement | ↔ | is provided as a guideline for accurate dosing of OPTIMMUNE.

**LEAFLET**

**PRESCRIPTION ANIMAL REMEDY  
KEEP OUT OF THE REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY**

**OPTIMMUNE ®OPHTHALMIC OINTMENT**

**ACTIVE CONSTITUENT:** 2 mg/g CYCLOSPORIN

For the treatment of idiopathic chronic keratoconjunctivitis sicca (KCS) and chronic superficial keratitis (CSK) in dogs.

**DIRECTIONS FOR USE****Dosage and administration**

Remove debris from affected eye(s) with suitable non-irritating solution.

For KCS apply a 6 mm strip of ointment to the affected eye(s) every 12 hours.

For CSK apply a 6-10 mm strip of ointment to the affected eye(s) every 12 hours.

Place the ointment directly on the cornea or into the conjunctival sac. Continue treatment while condition persists.

Dogs afflicted with KCS or CSK may require lifelong consistent therapy.

**General directions**

For best results in treating KCS the ointment should be administered early in the course of the disease before irreversible damage to the lacrymal tissue, or dense corneal scarring or pigmentation occurs. It is recommended that dogs exhibiting chronic recurring conjunctivitis be tested for adequate tear production.

The clinical effect of OPTIMMUNE Ophthalmic Ointment has not been determined in dogs with KCS due to the following conditions: congenital alacrima, sulfonamide usage, canine distemper virus, metabolic disease, surgical removal of the third eyelid gland and facial nerve paralysis with loss of the palpebral reflex.. Several days to a few weeks of application may be required before the clinical effects of Optimimmune are of sufficient magnitude to allow the safe withdrawal of previously initiated therapy.

Abrupt withdrawal of a therapeutic agent immediately upon initiation of treatment with Optimimmune can result in rapid clinical relapse.

**Contraindications/precautions**

Do not use if pre-existing viral or fungal ocular infections are present. Delay use until such infections have been successfully treated

The safety of Optimimmune in puppies, pregnant bitches, or dogs used for breeding has not been determined.

On rare occasions, instillation of OPTIMMUNE may be associated with stinging. If this persists beyond 7 days treatment should be withdrawn.

Additional information is in the Material Safety Data Sheet.

**Dispose** of empty container by wrapping with paper and putting in garbage.

**Store** below 25°C (Air conditioning).

Schering-Plough Animal Health  
(A Division of Schering-Plough Pty Ltd)  
11 Gibbon Road, Baulkham Hills NSW 2153

NRA 51551/0600