

Main panel – draft carton label



**PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**



**DEXASON ANTI-INFLAMMATORY AND
GLUCOGENIC STEROID INJECTION**

2mg/mL DEXAMETHASONE
(as SODIUM PHOSPHATE)

ILIUM 50mL

Ancillary panels

Anti-Inflammatory and Glucogenic Steroid Injection for Dogs, Cats, Horses, Cattle, Foals, Calves and Pigs.

DIRECTIONS FOR USE:

Dosage and Administration

Use the contents within 24 hours of first broaching of the vial. Discard the unused portion.

DOGS & CATS: 0.50-2mg, i.m. or i.v.

HORSES & CATTLE: 10-30mg, i.m. or i.v.

FOALS, CALVES & PIGS: 1mg/10kg, i.m. or i.v.

REPEAT EVERY 12 to 24 HOURS

WITHHOLDING PERIODS:

MILK: Milk collected from cows within 72 hours (6 milkings) following treatment **MUST NOT BE USED** for human consumption or processing. This milk should not be fed to bobby calves.

MEAT: **DO NOT USE** less than 10 days before slaughter for human consumption.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact the manufacturer on (02) 9604 6266 before using this product.

First Aid:

If poisoning occurs, contact a doctor or Poisons Information Centre.

Phone Australia 131126

TROY LABORATORIES PTY. LTD

31-35 Glendenning Road, Glendenning NSW 2761

website: www.troylab.com.au email: info@troylab.com.au

Dispose of empty container by wrapping with paper and putting in the garbage.

Store below 30°C (Room Temperature). Protect from light.

APVMA Approval No. 52859/50420

Batch:

Expiry:

Main panel – bottle label

**PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**



**DEXASON ANTI-INFLAMMATORY AND
GLUCOGENIC STEROID INJECTION**

2mg/mL DEXAMETHASONE
(as SODIUM PHOSPHATE)

ILIUM 50mL

Ancillary panel

Anti-Inflammatory and Glucogenic Steroid Injection for Dogs, Cats, Horses, Cattle, Foals, Calves and Pigs.

DIRECTIONS FOR USE:

Dosage and Administration

Use the contents within 24 hours of first broaching of the vial. Discard the unused portion.

DOGS & CATS: 0.50-2mg, i.m. or i.v.

HORSES & CATTLE: 10-30mg, i.m. or i.v.

FOALS, CALVES & PIGS: 1mg/10kg, i.m. or i.v.

REPEAT EVERY 12 to 24 HOURS

WITHHOLDING PERIODS:

MILK: Milk collected from cows within 72 hours (6 milkings) following treatment **MUST NOT BE USED** for human consumption or processing. This milk should not be fed to bobby calves.

MEAT: **DO NOT USE** less than 10 days before slaughter for human consumption.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): ESI Not established.

First Aid:

If poisoning occurs, contact a doctor or Poisons Information Centre.

Phone Australia 131126

TROY LABORATORIES PTY. LTD

31-35 Glendenning Road, Glendenning NSW 2761 Australia.

Dispose of empty container by wrapping with paper and putting in the garbage.

Store below 30°C (Room Temperature). Protect from light.

APVMA Approval No. 52859/50420

Batch:

Expiry: