



**Company Name:** INTERVET AUSTRALIA PTY LIMITED  
**Product Name:** NOBIVAC DHPPI CONTINUUM VACCINE

**APVMA Approval No:** 56912/100922



<b>Label Name:</b>	NOBIVAC DHPPI CONTINUUM VACCINE
<b>Signal Headings:</b>	PRESCRIPTION ANIMAL REMEDY KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY
<b>Constituent Statements:</b>	<b>ACTIVE CONSTITUENTS:</b> This vaccine contains freeze-dried living attenuated strains of canine distemper virus (CDV) strain Onderstepoort, canine adenovirus type 2 (CAV2) strain Manhattan LPV3 canine parvovirus (CPV) strain 154 and canine parainfluenza (CPi) strain Cornell grown in cell-line tissue culture. It is presented as a freeze-dried plug in single dose vials containing not less than $10^{4.0}$ TCID <sub>50</sub> CDV, $10^{4.0}$ TCID <sub>50</sub> CAV2, $10^{7.0}$ TCID <sub>50</sub> CPV and $10^{5.5}$ TCID <sub>50</sub> CPi attenuated vaccine virus. Each vial contains a single dose for reconstitution.
<b>Claims:</b>	Live vaccine to provide protection against canine distemper (CDV), infectious canine hepatitis caused by canine adenovirus type 1 (CAV1), canine parvovirus disease (CPV), respiratory disease caused by canine parainfluenza (CPi), canine adenovirus type 2 (CAV2) and to prevent viral excretion caused by canine parvovirus infection.
<b>Net Contents:</b>	single dose vial 10x single dose vials (1mL after reconstitution) 20x single dose vials (1mL after reconstitution) 25x single dose vials (1mL after reconstitution) 50x single dose vials (1mL after reconstitution)
<b>Directions for Use:</b>	<b>DIRECTIONS FOR USE</b> USE ALL PRODUCT WITHIN 30 MINUTES OF RECONSTITUTION.
<b>Precautions:</b>	<b>MANDATORY WARNINGS:</b> Only healthy dogs should be vaccinated and an adequate clinical examination should be made prior to inoculation. Sterile equipment should be used for administration, but avoid contamination of vaccine with traces of disinfectant or spirit.

<b>Side Effects:</b>	Occasionally a mild anaphylactic type hypersensitivity reaction may occur after vaccination, as is possible after inoculation of any foreign protein. Such types of reaction are in most cases self-limiting but may occasionally need veterinary intervention.
----------------------	---

<b>Dosage and Administration:</b>	<b>DOSAGE AND ADMINISTRATION:</b> The contents of one vial of reconstituted vaccine should be injected subcutaneously. Reconstitute immediately prior to use by the addition of 1.0 mL of sterile water for injection or sterile phosphate buffer.
-----------------------------------	---

<b>General Directions:</b>	<p><b>GENERAL DIRECTIONS:</b></p> <p>Vaccination may commence from 6 weeks of age. The final vaccination of the puppy course should be administered no earlier than 10 weeks of age provided at least 2 weeks has elapsed since the initial vaccination, with a booster 12 months later. In older puppies (10 weeks of age or older) only a single priming vaccination is necessary for protection against distemper, hepatitis and parvovirus with a booster 12 months later. For an optimal response to the parainfluenza component, animals should be vaccinated twice, 2 to 4 weeks apart, with the final vaccination at 10 weeks of age or older.</p> <p>It is recommended that dogs be revaccinated against distemper, hepatitis and parvovirus every 3 years. It is recommended that dogs be revaccinated for the parainfluenza component every 12 months – see vaccination program.</p> <p>Vaccination program:</p> <p>Puppies:</p> <p>Recommended program for pups where vaccination is started at 6-8 weeks:</p> <p>Age 6-8 weeks Nobivac® DHPPi Age 10 weeks Nobivac® DHPPi Where vaccination commences at 10 weeks or beyond: Nobivac® DHPPi followed by Nobivac® Pi 2 to 4 weeks later</p> <p>Booster vaccination</p> <p>1st annual booster Nobivac® DHPPi 2nd annual booster Nobivac® Pi 3rd annual booster Nobivac® Pi 4th annual booster Nobivac® DHPPi</p> <p>Adult Dogs:</p> <p>A single vaccination with Nobivac® DHPPi will provide at least three years protection against distemper, hepatitis and parvovirus and at least 12 months protection against parainfluenza.</p> <p>This product has been assessed as providing at least 36 months protection for distemper, hepatitis and parvovirus and 12 months protection for the parainfluenza component. Many factors influence the effectiveness of vaccination and the need for revaccination. The vaccination program for an individual animal should be determined within a veterinarian-client-patient relationship taking all these factors into account.</p> <p>Booster vaccination</p> <p>1st annual booster Nobivac® Pi 2nd annual booster Nobivac® Pi 3rd annual booster Nobivac® DHPPi 4th annual booster Nobivac® Pi</p> <p><b>FURTHER INFORMATION:</b></p> <p>Experience has shown that the maternal antibody status of pups within a litter varies greatly and reliance should not be placed on serological examination of the bitch alone.</p> <p><b>OVERDOSAGE:</b></p> <p>No particular signs at 10-fold overdose.</p>
----------------------------	--

<b>First Aid Instructions:</b>	<b>FIRST AID:</b> If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126
--------------------------------	---

<b>First Aid Warnings:</b>	
----------------------------	--

<b>Additional User Safety:</b>	Additional User Safety Take care to avoid self-injection.
--------------------------------	--

<b>Disposal:</b>	<b>DISPOSAL:</b> Discarded vials should immediately be placed in a designated and appropriately labelled 'biologicals' container. If this option is not available, bury the vial/container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and roots. Discarded needles should immediately be placed in a designated and appropriately labelled 'sharps' container. Dispose of outer packaging by wrapping with paper and putting in garbage
------------------	---

<b>Storage:</b>	Store between 2°C and 8°C (Refrigerate. Do not freeze). Protect from light.
-----------------	---