

APVMA FILE: 60441\_46408\_300Tab\_immediate\_MPL\_V02.pdf

Dimensions: label = 150mm (wide) x 100mm (high)

Scale: 100% of A4

Date: 28 July 2009

**PRESCRIPTION ANIMAL REMEDY**

KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY

## Cephaforte 600

### Antibiotic tablets for dogs

ACTIVE CONSTITUENT:  
Each tablet contains 600 mg CEPHALEXIN  
(as monohydrate)

For the treatment of infections in dogs caused by cephalixin sensitive organisms, including infections of soft cutaneous tissue, infections of the respiratory, urinary and digestive tracts, and for the prevention and treatment of osteomyelitis.

300 TABLETS

**Jurox**

**DIRECTIONS FOR USE:**

**Contraindication:** This product is contraindicated in animals with a history of allergic reactions to cephalosporins or other penicillins.  
Administer orally: Give 15 mg/kg bodyweight twice per day (equivalent to 1 Cephaforte 600 Tablet per 40 kg bodyweight given night and morning). Duration of treatment is at least 3 days, but may be extended at the discretion of the veterinarian. Nausea and vomiting in dogs may occasionally occur, but these effects are unusual and transitory.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

(B)

EXP.

407186

APVMA No. 60441/300 Tab/0309

JUROX Pty. Ltd.  
85 Gardiner Road  
Rutherford NSW 2320  
Infoline 1800 023 312  
Dispose of empty container  
by wrapping with paper and  
placing in garbage.  
Store below 30°C  
(Room Temperature)



---

## Primary container (carton) Main Panel

---

PRESCRIPTION ANIMAL REMEDY  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY



# CEPHAFORTE 600

## antibiotic tablets for dogs

Active Constituent:

Each tablet contains 600 mg CEPHALEXIN (as monohydrate)

**For the treatment of infections in dogs caused by cephalixin sensitive organisms, including infections of soft cutaneous tissue, infections of the respiratory, urinary and digestive tracts, and for the prevention and treatment of osteomyelitis.**

[60] [100] Tablets

[Jurox logo]

---

## Primary container (carton) Back Panel

---

### DIRECTIONS FOR USE

**Contraindication: This product is contraindicated for use in animals with a history of allergic reactions to cephalosporins or other penicillins.**

Administer orally. Give 15mg/kg bodyweight twice per day (equivalent to 1 Cephaforte 600 Tablet per 40kg bodyweight given night and morning).

Duration of treatment is at least 3 days, but may be extended at the discretion of the veterinarian.

Nausea and vomiting in dogs may occasionally occur, but these effects are unusual and transitory.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131126.

DISPOSE of empty container by wrapping with paper and placing in garbage.

[Jurox logo]

[Item code]

STORE below 30°C (Room Temperature)

Made in Australia by

Jurox Pty Ltd

85 Gardiner Rd

Rutherford NSW 2320

Infoline 1800 023 312

APVMA No. 60441/[10x6][10x10]Tab/0309

---

---

**Primary container (carton) Side Panel 1**

---

**CEPHAFORTE 600  
antibiotic tablets for dogs**



B.

EXP.

---

**Primary container (carton) Side Panel 2**

---

[BAR CODE]

---

**Primary container (carton) Side Tabs 1 and 2**

---

**CEPHAFORTE 600  
antibiotic tablets for dogs**

---

**Blister Label\***

---

FOR ANIMAL TREATMENT ONLY

**Cephaforte 600**

Antibiotic tablets for Dogs

600 mg cephalexin

[Jurox logo]

For oral use.

APVMA 60441/0309

[Item code]

B.

EXP.

---

*\*PLEASE NOTE: The blister pack is scored for subdivision. The batch and expiry date appear at least once on each blister strip. The other information appears in relation to at least every two dosage units.*

---