


Project: HRO7550	Description: Ilium Pentobarbitone 100mL carton	Material Type: SBB400	Product Code:	Date: 01/12/06
Dimensional Size: 55.00mm x 52.00mm x 100.00mm	Carton Area: 0.0357sq. m	Side Shown: printed		
Customer: TROY LABORATORIES	Operator: Justin Keane			
Style:	OZ ART ADVERTISING			
Overall Dimensions: 235 x 225.25				

Crease	-----



File Name: 61712_41426_100MLG_CARTON_MPL_V1
Dimensions: 100 x 55 x 52mm
Scale: A4 = 70% to scale
Date: 21st Sep 2007

140mm

PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

ilium
Tolfejec

**Anti-inflammatory Injection
for Cattle and Pigs**

ACTIVE CONSTITUENT: 40mg/mL Tolfenamic acid

Non steroidal anti-inflammatory analgesic -
antipyretic for use in cattle and pigs

ilium **100mL**

READ THE ENCLOSED LEAFLET BEFORE USING THIS PRODUCT

DIRECTIONS FOR USE
Restrains: DO NOT inject cattle other than into muscle tissue high on the side of the neck (see leaflet for diagram). Injection of product into muscles other than as described is likely to result in residues in meat above the MRL.

DOSAGE AND ADMINISTRATION
Use the contents of this vial within six months of initial broaching. Discard the unused portion.

PIGS Pneumonia: 1mL per 20kg bw I.M. Treatment may be repeated once only after 48 hours. Mastitis: 1mL per 10kg bw I.V.	CATTLE Mastitis: 1mL per 20kg bw I.M. Treatment may be repeated once only after 48 hours. Mastitis: 1mL per 10kg bw I.V.
--	--

APPROVED LABEL

WITHHOLDING PERIODS
MEAT: Cattle - 10 days IM injection, 4 days IV injection;
Pigs - 6 days.
MILK: Cattle - 12 hours.

TROY LABORATORIES PTY. LIMITED
98 Long Street, SMITHFIELD NSW 2164

Dispose of empty container by wrapping with paper and putting in garbage.
Store below 30°C (Room Temperature) away from direct sunlight.
APVMA Approval: 61712/100MLG/C707
Batch: _____
Expiry: _____

L0158

52mm

File Name: 61712_41426_100ML_LABEL_MPL_V1
Dimensions: 52 x 140mm
Scale: A4 = 100% to scale
Date: 21st September 2007




Pantone 289



Black

105mm


PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY



ilium
Tolfejec

Anti-inflammatory Injection for Cattle and Pigs

ACTIVE CONSTITUENT: 40mg/mL Tolfenamic acid



Pharmacological properties and pharmacokinetic particulars

Tolfenamic acid (N-(2-methyl-3-chlorophenyl) anthranilic acid) is a non-steroidal anti-inflammatory drug (NSAID) belonging to the fenamate group. Tolfenamic acid exerts anti-inflammatory, analgesic and antipyretic activities. The anti-inflammatory activity of tolfenamic acid is mainly due to an inhibition of cyclo-oxygenase and thus to a reduction of the synthesis of prostaglandins and thromboxanes, which are important inflammatory mediators.

INDICATIONS

For use by or under the direction of registered veterinarian as an aid in the treatment of pneumonia and acute mastitis in cattle and metritis-mastitis-agalactia in pigs.

RESTRAINTS

DO NOT inject cattle other than into muscle tissues high on the side of the neck (see diagram). Injection of product into muscles other than as described is likely to result in residues in meat above the MRL.

PRECAUTIONS

Repeat treatments or higher doses could result in residues above the MRL unless the label withholding period is extended. The prescribing Veterinarian would need to advise on an extended withholding period. Safe use of ILIUM TOLFEJEC Injection during pregnancy has not been established.

ilium

148.5mm

File Name: 61712_41426_LEAFLET_V1
 Dimensions: 105mmX148.5mm
 Scale: A4 = 100% to scale
 Date: 21st September 2007

105mm



DOSAGE AND ADMINISTRATION

Use the contents of this vial within six months of initial broaching. Discard the unused portion.



CATTLE

Pneumonia: 2 mg/kg (1mL per 20kg bw) by intramuscular injection high in the neck. Treatment may be repeated once only after 48 hours.
Mastitis: 4mg/kg (1mL per 10kg bw) as a single intravenous injection.



PIGS

Metritis-mastitis-agalactia: 2mg/kg (1mL per 20kg bw) as a single intramuscular injection.

Injection site



WITHHOLDING PERIODS

MEAT: Cattle by intramuscular administration: DO NOT USE less than 10 days before slaughter for human consumption.

Cattle by intravenous administration: DO NOT USE less than 4 days before slaughter for human consumption.

Pigs by intramuscular administration: DO NOT USE less than 6 days before slaughter for human consumption.

MILK: Cattle by Intramuscular or Intravenous administration: Milk collected from cows within 12 hours (1 milking) MUST NOT BE USED for human consumption or processing. This milk should not be fed to bobby calves.



TROY LABORATORIES PTY LIMITED
98 LONG ST, SMITHFIELD NSW 2164
W: www.troylab.com.au
E: info@troylab.com.au

Dispose of empty container by wrapping with paper and putting in the garbage.

Store below 30°C (Room Temperature). Protect from light.

APVMA Approval No.: 61712/0707

SU2562

ilium

148.5mm

